arXiv:1208.3183v1 [math.DS] 15 Aug 2012

Stability of the Rhomboidal Symmetric-Mass Orbit

Lennard Bakker[a], Skyler Simmons[a,*]

[a]*Department of Mathematics, Brigham Young University, 275 TMCB, Provo, UT 84602, USA*

Abstract

We extend the result of Yan (2012) to the rhomboidal problem with masses 1, m, 1, m for some $0 < m \leq 1$. A symmetry-reduction technique presented by Roberts (2007) enables us to compute the linear stability of the orbit by computing a single number. The orbit is shown to be linearly stable except for a small neighborhood of $m = 0$. Additionally, analysis of a suitable Poincaré section suggests numerically that the orbit is nonlinearly stable as well except for small m.

Keywords: n-body problem, binary collision, regularization, linear stability, Poincaré Section

---

1. Introduction

The Newtonian n-body problem involves determining the motion of n point masses in space. The bodies are mutually attracted by the gravitational law demonstrated by Newton. Specifically, if $\{q_1, q_2, ..., q_n\}$ represent the positions of the bodies in $\mathbb{R}^k$ ($k = 1, 2$, or 3) with masses $\{m_1, m_2, ..., m_n\}$ respectively, then their motion is governed by the system of differential equations

$$\ddot{q}_i = \sum_{i \neq j} \frac{m_i m_j (q_j - q_i)}{|q_i - q_j|^3}, \tag{1}$$

where the dot represents the derivative with respect to time.

Periodic orbits of the n-body problem are of particular interest. The most noteworthy recent example is the three-body figure-eight orbit discovered numerically

---

[*] Corresponding author

Email addresses: `bakker@math.byu.edu` (Lennard Bakker), `xinkaisen@gmail.com` (Skyler Simmons)

by Moore in [1], and whose existence was proven analytically by Chenciner and Montgomery in [2]. This orbit features three equal masses which all travel on the same curve in a periodic fashion. This orbit was shown to be linearly stable by Roberts in [3] by using the twelve-fold symmetry present in the orbit, reducing the question of stability to a simple numerical calculation. Robert's technique has since been applied to other symmetric periodic orbits in various n-body problems (for instance [4], [5], [6], [7], and [8]).

Another important aspect of the n-body problem involves the study of singularities. A *collision singularity* is said to occur when $q_i = q_j$ for some $i \neq j$. Mathematically, this yields a zero denominator in (1). Certain classes of collision singularities are known to be *regularizable*, meaning that under a suitable change of space and time variables the orbit can be analytically continued past the collision time. In particular, binary collisions (where exactly two bodies collide) can be regularized. It is also possible to regularize simultaneous binary collisions (where multiple pairs of bodies collide at the same time) in certain cases when symmetries are present in the orbit. In [9] and [10], Sweatman analyzes a symmetric periodic orbit with both binary and simultaneous binary collisions in the collinear four-body problem. This orbit was proven to exist analytically by Ouyang and Yan in [11]. Later, a wider variety of periodic collision-based orbits was shown to exist by Shibayama in [12]. The key feature in each of these orbits is that they can be reduced to two degrees of freedom.

In this paper, we study a periodic orbit in the planar rhomboidal four-body problem, featuring binary collisions between alternating pairs of masses (as shown in Figure 1). This orbit was studied by Waldvogel in [13], who numerically studied the chaotic behaviors of the bodies, including escape, quasiperiodic orbits, resonances, and quadruple collision. The stability technique of Roberts was used in [7] to show that this orbit was linearly stable in the case where all masses are equal. Our work extends the stability calculations to the case where there are two pairs of equal masses. Under an appropriate rescaling, we need only consider a bounded interval of mass values, which simplifies the numerical calculations.

The remainder of the paper is as follows. In Section 2, we will describe the periodic orbit under consideration, and give the coordinate transformation which regularizes the binary collisions. Section 3 will outline the standard technique for showing linear stability for the orbit, as well as give the result of the symmetry-reduction technique developed by Roberts. Section 4 is divided into three portions



Figure 1: The rhomboidal four-body orbit.

detailing the numerical calculations performed. In particular, the first subsection describes how initial conditions for the orbit were obtained, the second gives the results for the numerical linear stability computation, and the third describes a Poincaré section analysis of the orbit.

2. The Rhomboidal Symmetric-Mass Orbit

We consider the planar Newtonian 4-body problem with bodies located at

$$q_1 = (x_1, 0), q_2 = (0, x_2), q_3 = -q_1, q_4 = -q_2 \tag{2}$$

and masses 1, m, 1, m respectively for some $m \in (0, 1]$. The bodies travel along the x and y axes, forming the vertices of a rhombus at all times away from collision. Binary collisions occur between the bodies with equal masses at the origin. For the periodic orbit, the non-colliding bodies have zero momentum at collision time. Following collision, the colliding bodies eject along the appropriate coordinate axis, and the remaining two bodies travel toward collision at the origin, where a similar (zero momentum of non-colliding bodies) behavior occurs. (See Figure 1.)

Analytic existence of this orbit was shown in [7] in the $m = 1$ case, and for

$m \in (0,1]$ in [12]. The Hamiltonian for the system is given by

$$H = \frac{1}{4}w_1^2 + \frac{1}{4m}w_2^2 - \frac{1}{2x_1} - \frac{m^2}{2x_2} - \frac{4m}{\sqrt{x_1^2 + x_2^2}}$$

where $w_1 = 2\dot{x}_1, w_2 = 2m\dot{x}_2$. We can continue the orbits past collision via a regularization under which binary collision corresponds to an elastic bounce. To regularize these collisions, we use a Levi-Civita-type change of coordinates. Using the canonical transformations $Q_i^2 = x_i$, $P_i = 2Q_i w_i$ for $i = 1,2$, with a change of time satisfying $dt/ds = x_1 x_2$, the regularized Hamiltonian in extended phase space is given by

$$\begin{aligned}
\Gamma &= \Gamma(Q_1, Q_2, P_1, P_2, E) \\
&= \frac{dt}{ds}(H - E) \\
&= \frac{1}{16}Q_2^2 P_1^2 + \frac{1}{16}\frac{Q_1^2 P_2^2}{m} - \frac{1}{2}Q_1^2 m^2 - \frac{1}{2}Q_2^2 - \frac{4Q_1^2 Q_2^2 m}{\sqrt{Q_1^4 + Q_2^4}} - Q_1^2 Q_2^2 E.
\end{aligned}$$

This yields the equations of motion

$$Q_1' = \frac{1}{8}Q_2^2 P_1, \tag{3}$$

$$Q_2' = \frac{1}{8m}Q_1^2 P_2, \tag{4}$$

$$P_1' = -\frac{1}{8m}Q_1 P_2^2 + Q_1 m^2 + \frac{8Q_1 Q_2^2 m}{(Q_1^4 + Q_2^4)^{1/2}} - \frac{8Q_1^5 Q_2^2 m}{(Q_1^4 + Q_2^4)^{3/2}} + 2Q_1 Q_2^2 E, \tag{5}$$

$$P_2' = -\frac{1}{8}Q_2 P_1^2 + Q_2 + \frac{8Q_1^2 Q_2 m}{(Q_1^4 + Q_2^4)^{1/2}} - \frac{8Q_1^2 Q_2^5 m}{(Q_1^4 + Q_2^4)^{3/2}} + 2Q_1^2 Q_2 E, \tag{6}$$

where $'$ denotes the derivative with respect to the new time variable s.

At the time of collision of the two bodies on the x-axis, we have $Q_1 = 0$ and $P_2 = 0$. At this time, setting $\Gamma = 0$ yields

$$Q_2^2\left(\frac{1}{16}P_1^2 - \frac{1}{2}\right) = 0.$$

Hence, $P_1 = \pm 8^{3/2}$, with the sign being the same as the sign on Q_2. Similarly, at the time of collision of the two bodies on the y-axis, $Q_2 = P_1 = 0$. The condition

$\Gamma = 0$ then gives

$$Q_1^2\left(\frac{1}{16m}P_2^2 - \frac{1}{2}m^2\right) = 0,$$

and so $P_2 = \pm(8m)^{3/2}$.

The symmetries present in the rhomboidal four-body orbit are the same as those presented in the collinear four-body orbit presented in Section 3 of [6]. Let

$$S = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}. \tag{7}$$

If $\gamma(s)$ is a T-periodic solution to (3) - (6), both $S\gamma(-s+T)$ and $-S\gamma(-s+T/2)$ are solutions as well. Existence and uniqueness of solutions then imply that

$$\gamma(-s+T) = S\gamma(s) \text{ and } \gamma(-s+T/2) = -S\gamma(s)$$

for all s. Hence the symmetry group for the rhomboidal four-body orbit is isomorphic to the Klein four group.

3. Linear Stability

Note that Γ is a smooth function defined on $\mathbb{R}^4 \setminus \{Q_1 = Q_2 = 0\}$. Suppose $\gamma(s)$ is a T-periodic solution of the system $z' = JD\Gamma(z)$, where $' = d/ds$,

$$J = \begin{bmatrix} O & I \\ -I & O \end{bmatrix},$$

and I and O are the 2×2 identity and zero matrices, respectively. If $X(s)$ is the fundamental matrix solution of the linearized equations

$$\xi' = JD^2(\gamma(s))\xi, \quad \xi(0) = I \tag{8}$$

then the monodromy matrix is given by $X(T)$ and satisfies $X(s+T) = X(s)X(T)$ for all s. Eigenvalues of the monodromy matrix are also the characteristic multipliers of γ, and therefore determine the linear stability of γ. In particular, γ is spectrally stable if all of its characteristic multipliers lie on the unit circle, and γ

is linearly stable if it is spectrally stable and semisimple apart from trivial eigenvalues.

It is also possible to find the characteristic multipliers of γ by using another initial condition $\xi(0)$ in (8). If Y_0 is an invertible matrix, let $Y(s)$ be the fundamental matrix solution to

$$\xi' = JD^2(\gamma(s))\xi, \quad \xi(0) = Y_0. \tag{9}$$

Then by definition of $X(s)$, we have $Y(s) = X(s)Y_0$, and so $X(T) = Y(T)Y_0^{-1}$. Then we have

$$X(T) = Y(T)Y_0^{-1} = Y_0(Y_0^{-1}Y(T)))Y_0^{-1}.$$

Hence, $Y_0^{-1}Y(T)$ is similar to $X(T)$, so the eigenvalues of $Y_0^{-1}Y(T)$ are the same as the characteristic multipliers of γ.

In [3], a technique for factoring the monodromy matrix in terms of symmetries of a periodic orbit is presented. These allow the linear stability of a periodic orbit to be determined by computing the eigenvalues of the fundamental matrix solution at time $s = T/N$ for some natural number N determined by the symmetry group of the orbit. Moreover, Roberts in [3] shows that if W is a symplectic matrix such that

$$\frac{1}{2}(W + W^{-1}) = \begin{bmatrix} K^T & O \\ O & K \end{bmatrix} \tag{10}$$

for some matrix K, then the eigenvalues of W lie on the unit circle if and only if all the eigenvalues of K are real and lie in the interval $[-1,1]$. Further, the eigenvalues W lie on the unit circle if and only if the eigenvalues of W^k lie on the unit circle for all $k \in \mathbb{Z}$. Each of these results combine to give the following linear stability result.

Theorem 1. *For the rhomboidal symmetric-mass orbit, there exists a choice of Y_0 such that $Y_0^{-1}Y(T) = W^2$ for a symplectic matrix W which satisfies (10). Moreover, K has the form*

$$K = \begin{bmatrix} -1 & * \\ 0 & c_2^T(SJc_4) \end{bmatrix},$$

where c_i represents the ith column of $Y(T/4)$. Therefore, the rhomboidal four-body orbit is linearly stable if and only if $c_2^T(SJc_4)$ is real and lies within the interval $[-1,1]$, with S as defined in (7).

The proof of this theorem is precisely sections 3.1 and 3.2 of [6], with greater detail presented in [3]. Although the arguments in that section are applied to a different orbit, both orbits have the same time-reversing symmetries, and the symmetries of the orbit are the only feature required for the arguments to hold.

4. Numerical Calculations

4.1. Determining Initial Conditions

In order to determine the initial conditions for the rhomboidal orbit, we model each of Q_1, Q_2, P_1, and P_2 by truncated trigonometric polynomials:

$$\tilde{Q}_1 = \sum_{i=0}^{n} a_i \sin((2i+1)s), \tag{11}$$

$$\tilde{Q}_2 = \sum_{i=0}^{n} b_i \sin((2i+1)(s+\pi/2)), \tag{12}$$

$$\tilde{P}_1 = \sum_{i=0}^{n} c_i \sin((2i+1)(s-\pi/2)), \tag{13}$$

$$\tilde{P}_2 = \sum_{i=0}^{n} d_i \sin((2i+1)s). \tag{14}$$

The choice of trigonometric polynomials is natural for modeling periodic behavior. A similar technique was carried out by Simó in [14]. The time shifts and choice of odd-only multiples of s correspond to symmetries of the orbit. In particular, for these polynomials, the time-reversing symmetries shown earlier are built-in, and the non-colliding bodies have zero net momentum at collision time. For a fixed n, we numerically minimize the value of

$$\int_0^{2\pi} \left((Q_1' - \tilde{Q}_1')^2 + (Q_2' - \tilde{Q}_2')^2 + (P_1' - \tilde{P}_1')^2 + (P_2' - \tilde{P}_2')^2 \right) \, ds$$

where the minimization is taken over the space of coefficients $\{a_i, b_i, c_i, d_i\}$. We combine this with a root-finding technique to find the appropriate value of E for a 2π-periodic orbit, as in [5]. Once these trigonometric polynomials are determined, we can extract the initial conditions for the orbit by evaluating $\tilde{Q}_1, \tilde{Q}_2, \tilde{P}_1, \tilde{P}_2$ at any fixed time $s \in [0, 2\pi]$.



Figure 2: The value of Q_2 as a function of m.



Figure 3: The value of E as a function of m.

We obtain the initial conditions for the 2π-periodic orbit for $m = 1$ by rescaling the conditions given in [7]. It is easy to check that if $(Q_1(s), Q_2(s), P_1(s), P_2(s), E)$ is a solution for (3) - (6), then $(\varepsilon Q_1(\varepsilon s), \varepsilon Q_2(\varepsilon s), P_1(\varepsilon s), P_2(\varepsilon s), E/\varepsilon^2)$ also satisfies (3) - (6). Moreover, this rescaling does not change the linear stability of the orbit. Given the initial conditions in [7] and rescaling with $\varepsilon \approx 1.55$, the orbit is roughly 2π-periodic, which we verify by integration using the standard fourth-order Runge-Kutta-Fehlberg algorithm. A standard curve-fitting technique can then be used to give the coefficients $\{a_i, b_i, c_i, d_i\}$ in the equations (11) - (14). After that, a gradual "step-down" technique (as in [5]) can be used to find the initial conditions for other values of $m \in (0, 1]$. If we assume the initial conditions occur at the time the two bodies with mass 1 collide, we have $Q_1 = P_2 = 0$. Also, as before, we know that $P_1 = -2\sqrt{2}$, so we need only to find the values of Q_2 and E. The results of the numerical calculation are shown in Figures 2 - 5.

4.2. *Linear Stability Computations*

Using the initial conditions for $\gamma(s)$ leading to a 2π-periodic orbit given earlier, we numerically compute $Y(T/4)$ with the initial condition

$$Y_0 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \end{bmatrix}$$



Figure 4: Numerical integrations of (3) - (6) with initial conditions obtained from the trigonometric polynomials $\tilde{Q}_i, \tilde{P}_i$ for various masses.



Figure 5: Orbits in terms of the original physical variables x_1 and x_2 obtained by coordinate transformation of the data shown in Figure 4.

from Theorem 1 using a standard Runge-Kutta-Fehlberg algorithm with a fixed time step

$$\Delta t = \frac{\pi/2}{500000} = \frac{\pi}{1000000}.$$

The value of $c_2^T(SJc_4)$ plotted against m is given in Figure 6. Numerically, the computed values of $c_2^T(SJc_4)$ lie below 1 for all $m \geq .04$, and above 1 for $m \leq .03$. Accounting for numerical error, we have shown the following

Theorem 2. *There exists some positive number* ε *such that the symmetric mass periodic orbit of the regularized planar rhomboidal four-body problem (3) - (6) is linearly stable for* $m \in (.03+\varepsilon, 1]$.

Note that in the case $m = 1$, our computation confirms the calculation given in [7], wherein an additional symmetry is present in the orbit.



Figure 6: The value of $c_2^T(SJc_4)$ as a function of m.

4.3. *Poincaré Section Analysis*

To numerically analyze nonlinear stability, we find a suitable Poincaré section for the orbit. This was done in the $m = 1$ case in [13]. Our more general Poincaré section is based on techniques presented in [15] and [9]. For any value of m, we seek a number α such that

$$\frac{x_1}{x_2} = \alpha$$

is maintained throughout the orbit, with x_1 and x_2 as defined in (2) earlier. In other words, the value of α corresponds to the ratio of x_1 and x_2 in a homographic orbit where the trajectories of the bodies correspond to total collapse (or ejection from total collapse). We find this value of α by solving the standard equations of motion (1) with the substitutions $x_1 = \alpha x_2$ and $\ddot{x_1} = \alpha \ddot{x_2}$. Doing so, we find that the required value of α for a given mass m is the single positive root of the 12th-degree polynomial

$$(1+\alpha^2)^3(1-m\alpha^3)^2 - 64\alpha^6(m-1)^2 = 0.$$

The value of α as a function of m is plotted in Figure 7.



Figure 7: The value of α as a function of m.

For fixed $E = -1$, we define a Poincaré section Σ to be the two-dimensional surface given by $x_1 = \alpha x_2$ in the phase space defined by the variables $(x_1, x_2, \dot{x}_1, \dot{x}_2)$. (Note that $\dot{x}_1$ and $\dot{x}_2$ are simply linear re-scalings of w_1 and w_2.) Restricting to $E = -1$, we find a bound on the possible values of x_1. Specifically, if $\dot{x_1} = \dot{x_2} = 0$ on Σ, the condition $E = -1$ requires that

$$x_1 = \frac{1}{2} + \frac{m^2\alpha}{2} + \frac{4m\alpha}{\sqrt{\alpha^2+1}} = r_{\max}.$$

For a set of initial conditions on Σ, the requirement $E = -1$ necessarily implies that $x_1 \leq r_{\max}$, and if either of $\dot{x}_1$ or $\dot{x}_2$ are non-zero, then the strict inequality $x_1 < r_{\max}$ holds.

We define coordinates (r, θ) on Σ by

$$r = \frac{x_1}{r_{\max}}, \quad \theta = \tan^{-1}\left(\frac{\dot{x}_1}{\alpha \dot{x}_2}\right).$$

Under this change of coordinates, the homographic orbit corresponds to the line $\theta = \pi/4$. For a 9×15 grid of equally spaced initial conditions in (r, θ) we numerically integrate (3) - (6) for the corresponding initial conditions and record the first 200 intersections of the orbit with Σ. (Integration was preemptively terminated if any of Q_i, P_i exceeded 1000 in absolute value.) The results of this are shown in Figures 8 - 12. The observed concentric rings numerically match the predicted result of Moser's Invariant Curve Theorem in [16], and show that the rhomboidal symmetric-mass orbit is nonlinearly stable for $m \in (.03 + \varepsilon, 1]$ for the same ε as in Theorem 2.



Figure 8: Poincaré sections plotted for $m = .1$ (top) and $m = .2$ (bottom). In these plots, r lies on the vertical axis. The homographic orbit at $\theta = \pi/4$ is not plotted for clarity.



Figure 9: Poincaré sections plotted for $m = .3$ (top) and $m = .4$ (bottom).



Figure 10: Poincaré sections plotted for $m = .5$ (top) and $m = .6$ (bottom).



Figure 11: Poincaré sections plotted for $m = .7$ (top) and $m = .8$ (bottom).



Figure 12: Poincaré sections plotted for $m = .9$ (top) and $m = 1$ (bottom).

References

[1] C. Moore, Braids in classical dynamics, Phys. Rev. Lett. 70 (1993) 3675–3679.

[2] A. Chenciner, R. Montgomery, A remarkable periodic solution of the three-body problem in the case of equal masses, Ann. of Math. (2) 152 (2000) 881–901.

[3] G. E. Roberts, Linear stability analysis of the figure-eight orbit in the three-body problem, Ergodic Theory Dynam. Systems 27 (2007) 1947–1963.

[4] L. F. Bakker, S. Mancuso, S. C. Simmons, Linear stability for some symmetric periodic simultaneous binary collision orbits in the planar pairwise symmetric four-body problem, J. Math. Anal. Appl. 392 (2012) 136–147.

[5] L. F. Bakker, T. Ouyang, D. Yan, S. Simmons, Existence and stability of symmetric periodic simultaneous binary collision orbits in the planar pairwise symmetric four-body problem, Celestial Mech. Dynam. Astronom. 110 (2011) 271–290.

[6] L. F. Bakker, T. Ouyang, D. Yan, S. Simmons, G. E. Roberts, Linear stability for some symmetric periodic simultaneous binary collision orbits in the four-body problem, J. Math. Anal. Appl. 108 (2010) 147–164.

[7] D. Yan, Existence and linear stability of the rhomboidal periodic orbit in the planar equal mass four-body problem, J. Math. Anal. Appl. 388 (2012) 942–951.

[8] D. Yan, Existence of the Broucke periodic orbit and its linear stability, J. Math. Anal. Appl. 389 (2012) 656–664.

[9] W. L. Sweatman, The symmetrical one-dimensional Newtonian four-body problem: a numerical investigation, Celestial Mech. Dynam. Astronom. 82 (2002) 179–201. The restless universe (Blair Atholl, 2000).

[10] W. L. Sweatman, A family of symmetrical Schubart-like interplay orbits and their stability in the one-dimensional four-body problem, Celestial Mech. Dynam. Astronom. 94 (2006) 37–65.

[11] T. Ouyang, D. Yan, Periodic solutions with alternating singularities in the collinear four-body problem, Celestial Mech. Dynam. Astronom. 109 (2011) 229–239.

[12] M. Shibayama, Minimizing periodic orbits with regularizable collisions in the n-body problem, Arch. Ration. Mech. Anal. 199 (2011) 821–841.

[13] J. Waldvogel, The rhomboidal symmetric four-body problem, Celestial Mech. Dynam. Astronom. 113 (2012) 113–123.

[14] C. Simó, New families of solutions in N-body problems, in: European Congress of Mathematics, Vol. I (Barcelona, 2000), volume 201 of *Progr. Math.*, Birkhäuser, Basel, 2001, pp. 101–115.

[15] J. Hietarinta, S. Mikkola, Chaos in the one-dimensional gravitational three-body problem, Chaos 3 (1993) 183–203.

[16] C. L. Siegel, J. K. Moser, Lectures on celestial mechanics, Classics in Mathematics, Springer-Verlag, Berlin, 1995.